

Temporal dynamic of strongly coupled epsilon-near-zero media

**Mehdi H. Ebrahim¹, Andrea Marini², Nathaniel Kinsey³, Jacob B. Khurgin⁴, Lucia Caspani⁵,
Marcello Ferrera⁶, Vladimir M. Shalaev⁷, Alexandra Boltasseva⁷, Daniele Faccio⁸, Matteo Clerici¹**

¹James Watt School of Engineering, University of Glasgow, G12 8QQ Glasgow, UK

²Department of Physical and Chemical Sciences, University of L'Aquila, Via Vetoio, 67100 L'Aquila, Italy

³Department of Electrical and Computer Engineering, Virginia Commonwealth University, Richmond, Virginia 23284, USA

⁴Department of Electrical and Computer Engineering, Johns Hopkins University, Baltimore, Maryland 21218, USA

⁵Institute of Photonics, Department of Physics, University of Strathclyde, G1 1RD, Glasgow, UK

⁶Institute of Photonics and Quantum Sciences, Heriot-Watt University, SUPA, Edinburgh EH14 4AS, UK

⁷School of Electrical and Computer Engineering, Purdue University, West Lafayette, Indiana 47907, USA

⁸School of Physics and Astronomy, University of Glasgow, G12 8QQ Glasgow, UK

*corresponding author, E-mail: matteo.clerici@glasgow.ac.uk

Abstract: We shall present an overview of our recent investigations into the time-dependent effects observed in transparent conductive oxides photoexcited at the epsilon near zero point. We will then discuss the role of the epsilon-near-zero mode in establishing the temporal dynamics in strongly coupled plasmonic systems.